



```

5'-TGTGGAAAGGACGAGGATCC[...shRNA oligo cloning site...]GAATTCTACCGGTAGGGGAGGCGCTTTTCCCAAGGCAGT-3'
           BamH I                         EcoR I
3'-ACACCTTCTCTGCTCTAGG[...shRNA oligo cloning site...]CTTAAGATGGCCATCCCTCCGCGAAAAGGGTTCCGTCA-5'

```

**Restriction Map and Cloning Site of RNAi-Ready pSIREN-RetroQ-TetH Retroviral Vector.** All restriction sites shown are unique. RNAi-Ready pSIREN-RetroQ-TetH is provided as a linearized vector digested with *BamH I* and *EcoR I*. Nucleotides in gray were removed during linearization. This linearized vector is ready for ligation of an appropriate shRNA oligo containing *BamH I* and *EcoR I* overhangs.

**Description**

RNAi-Ready pSIREN-RetroQ-TetH is a self-inactivating retroviral expression vector designed to express a ds short hairpin RNA (shRNA) under the control of the modified Tet-responsive promoter ( $P_{TREM_{mod}/U6}$ ) derived from the  $P_{TREM_{mod}}$  and the human U6 promoter ( $P_{U6}$ ). RNAi-Ready pSIREN-RetroQ-TetH is provided as a linearized vector digested with *BamH I* and *EcoR I*. It is used for targeted and inducible gene silencing when a DNA oligonucleotide encoding an appropriate shRNA is ligated into the vector. shRNA expression is controlled by the tetracycline suppressor tTS (1).  $P_{TREM_{mod}}$  consists of seven direct repeats of a 36 bp sequence that contains the 19 bp tet operator sequence (tetO). You can transfect your pSIREN-RetroQ-TetH construct as a plasmid expression vector, or—upon transfection into a packaging cell line—this vector can transiently express, or integrate and stably express a viral genomic transcript containing the  $P_{TREM_{mod}/U6}$  promoter and the shRNA. The vector contains a hygromycin resistance gene (*Hyg<sup>r</sup>*) under the control of the murine phosphoglycerate kinase (*PGK*) promoter ( $P_{PGK}$ ) for the selection of stable transfectants.

This retroviral vector is optimized to eliminate promoter interference through self-inactivation. The hybrid 5' LTR consists of the cytomegalovirus (CMV) type I enhancer and the mouse sarcoma virus (MSV) promoter. This construct drives high levels of transcription in HEK 293-based packaging cell lines due, in part, to the presence of adenoviral E1A (2–5) in these cells. The self-inactivating feature of the vector is provided by a deletion in the 3' LTR enhancer region (U3). During reverse transcription of the retroviral RNA, the inactivated 3' LTR is copied and replaces the 5' LTR, resulting in inactivation of the 5' LTR CMV enhancer sequences. This mechanism may reduce the phenomenon known as promoter interference (6, 7) and allow more efficient expression. Viral infection of host cells with recombinant pSIREN-RetroQ-TetH is the preferred delivery method.

(PR611409; published 13 February 2006)



Clontech

United States/Canada  
800.662.2566  
Asia Pacific  
+1.650.919.7300  
Europe  
+33.(0)1.3904.6880  
Japan  
+81.(0)77.543.6116

Clontech Laboratories, Inc.  
A Takara Bio Company  
1290 Terra Bella Ave.  
Mountain View, CA 94043  
Technical Support (US)  
E-mail: tech@clontech.com  
www.clontech.com

Also included in the viral genomic transcript are the necessary viral RNA processing elements including the LTRs, packaging signal ( $\Psi^+$ ), and tRNA primer binding site. RNAi-Ready pSIREN-RetroQ-TetH also contains a bacterial origin of replication and *E. coli* Amp<sup>r</sup> gene for propagation and selection in bacteria.

## Use

RNAi-Ready pSIREN-RetroQ-TetH is used for targeted and inducible gene silencing when a DNA oligonucleotide encoding an appropriate shRNA is inserted. To construct recombinant pSIREN-RetroQ-TetH-shRNA constructs, first design, generate, and anneal complementary shRNA oligonucleotides using the protocols in the Knockout Inducible RNAi Systems User Manual (PT3810-1). The annealed oligonucleotide should contain 5'-*Bam*H I and 3'-*Eco*R I sites. Then ligate the annealed oligonucleotide into the RNAi-Ready pSIREN-RetroQ-TetH vector.

Your pSIREN-RetroQ-TetH construct can be transfected as a plasmid expression vector to screen for functional shRNA oligonucleotides. For gene silencing experiments using viral delivery, transfect the pSIREN-RetroQ-TetH construct into a packaging cell line (such as the Retro-X™ Universal Packaging System [Cat. No. 631512]). RNA from the vector is packaged into infectious retroviral particles. These infectious particles are replication-incompetent since pSIREN-RetroQ-TetH lacks structural genes (*gag*, *pol*, and *env*) necessary for particle formation and replication; however, these genes are stably integrated as part of the packaging cell genome. These retroviral particles can infect a wide range of target cells and transmit the shRNA but cannot replicate within these cells due to the absence of viral structural genes. The separate introduction and integration of the structural genes into the packaging cell line minimizes the chances of producing replication-competent virus due to recombination events during cell proliferation.

**pSIREN-RetroQ-TetH-Luc Vector** contains a validated luciferase shRNA oligonucleotide insert. When tested in HEK 293 cells plated at  $1 \times 10^5$  per well and transfected with a 2:1:1 ratio of ptTS-Neo:pSIREN-RetroQ-TetH-Luc:pCMV-Luc, we consistently observe a 60–75% knockdown of luciferase activity with 48 hr of 1  $\mu$ g/ml Dox induction. This control vector allows you to directly monitor the cloning efficiency of your shRNA insert into the pSIREN-RetroQ-TetH Vector.

## Location of Features

- PGK promoter ( $P_{PGK}$ ): 2–510
- Hygromycin resistance gene (*Hyg<sup>r</sup>*): 528–1495
- 3' MoMuLV LTR (deletion in U3): 2164–2589  
Poly A region: 2415–2430
- SV40 promoter: 2869–3155
- SV40 ori: 3090–3156
- Col E1 ori (site of replication initiation): 4074–3475
- Ampicillin resistance gene ( $\beta$ -lactamase): 5096–4236  
Start codon (ATG): 5096–5094      Stop codon (TAA): 4238–4236
- 5' LTR (CMV/MSV): 5459–6186  
Cytomegalovirus (CMV)/ mouse sarcoma virus (MSV) hybrid promoter: 5459–5969  
R region: 6042–6112  
U5 region: 6114–6186
- $\Psi^+$  (extended packaging signal): 6216–7025
- $P_{tight}$  Tet-responsive promoter: 7041–7321  
Tet responsive element ( $TRE_{mod}$ ): 7041–7304  
Location of seven tetO 19-mers: 7064–7082 ; 7100–7018 ; 7135–7153 ; 7171–7189 ; 7207–7225; 7242–7250  
& 7278–7296
- U6 promoter ( $P_{U6}$ ): 7323–7389
- TATA box: 7367–7372

### Sequencing Primer Location

- U6 Forward Sequencing Primer: 6958-6985  
5'-CTTGAACCTCCTCGTTTCGACCCCGCCTC-3'

### Selection of Stable Transfectants

- Selectable marker: plasmid confers resistance to hygromycin.

### Propagation in *E. coli*

- Suitable host strains: DH5 $\alpha$ , DH10B, and other general purpose strains.
- Selectable marker: plasmid confers resistance to ampicillin (100  $\mu$ g/ml) in *E. coli* hosts.
- *E. coli* replication origin: Col E1
- Copy number: low

### References

1. Freundlieb S., Schirra-Muller, C. & Bujard, H. (1999) *J. Gene Med.* **1**(1):4–12.
2. Kinsella, T. M. & Nolan G. P. (1996) *Hum. Gene Ther.* **7**(18):1405–1413.
3. Ory, D. S., Neugeboren, B. A. & Mulligan, R. C. (1996) *Proc. Nat. Acad. Sci. USA* **93**(21):11400–11406.
4. Pear, W. S., Nolan, G. P., Scott, M. L. & Baltimore, D. (1993) *Proc. Natl. Acad. Sci. USA* **90**(18):8392–8396.
5. Yang, S., Delgado, R., King, S. R., Woffendin, C., Barker, C. S., Yang, Z. Y., Xu, L., Nolan, G. P. & Nabel, G. J. (1999) *Hum. Gene Ther.* **10**(1):123–132.
6. Barton, G.M. & Medzhitov R. (2002) *Proc. Natl. Acad. Sci. USA* **99**(23):14943–14945.
7. Emerman, M. & Temin, H. M. (1984) *Cell* **39**(3 pt. 2):449–467.

**Note:** The attached sequence file has been compiled from information in the sequence databases, published literature, and other sources, together with partial sequences obtained by Clontech. This vector has not been completely sequenced.

The viral supernatants produced by this retroviral vector could, depending on your cloned insert, contain potentially hazardous recombinant virus. Due caution must be exercised in the production and handling of recombinant retrovirus. Appropriate NIH, regional, and institutional guidelines apply.

### Notice to Purchaser

This product is intended to be used for research purposes only. It is not to be used for drug or diagnostic purposes nor is it intended for human use. Clontech products may not be resold, modified for resale, or used to manufacture commercial products without written approval of Clontech Laboratories, Inc.

Use of the Tetracycline controllable expression systems (the "Tet Technology") is covered by a series of patents including U.S. patent Nos. 5,464,758 and 5,814,618, which is proprietary to TET Systems Holding GmbH & Co. KG. Academic research institutions are granted an automatic license with the purchase of this product to use the Tet Technology only for internal, academic research purposes, which license specifically excludes the right to sell, or otherwise transfer, the Tet Technology or its component parts to third parties. In accepting this license, all users acknowledge that the Tet Technology is experimental in nature. TET Systems Holding GmbH & Co. KG makes no warranties, express or implied or of any kind, and hereby disclaims any warranties, representations, or guarantees of any kind as to the Tet Technology, patents, or products. All others are invited to request a license from TET Systems Holding GmbH & Co. KG prior to purchasing these reagents or using them for any purpose. Clontech is required by its licensing agreement to submit a report of all purchasers of the Tet-controllable expression system to IP Merchandisers, Inc. For license information, please contact:

Hans Peter Kneubuehl  
TET Systems Holding GmbH & Co. KG  
Im Neuenheimer Feld 582  
69120 Heidelberg, Germany  
Tel +49 6221 588 04 00  
Fax +49 6221 588 04 04  
eMail: [kneubuehl@tet-systems.de](mailto:kneubuehl@tet-systems.de)  
or use our electronic licensing request form via <http://www.tetsystems.com/licensing/index.html>

This product is sold under license from the Fred Hutchinson Cancer Research Center. Rights to use this product are limited to research only. No other rights are conveyed. Inquiry into the availability of a license to broader rights or the use of this product for commercial purposes should be directed to Fred Hutchinson Cancer Research Center, Technology Transfer Office, 1100 Fairview Avenue North, C2M-027, Seattle, WA 98109. Purchase of this product does not grant rights to: (1) offer the materials or any derivatives thereof for resale; or (2) to distribute or transfer the materials or any derivatives thereof to third parties.

Clontech, Clontech Logo and all other trademarks are the property of Clontech Laboratories, Inc. Clontech is a Takara Bio Company. ©2006