



Certificate of Analysis

rLV.EF1.tdTomato-9 0036VCT / 0008VCT

rV2.1A1.5764 C2

rLV: lentiviral vector/ EF1: Elongation Factor 1 alpha promoter / tdTomato

User manuals are available at: https://www.takarabio.com/.

Passed

BIOLOGICAL PRODUCTS: Defective lentiviral based vectors expressing a sequence of interest

Country of Origin: FRANCE

Result Unit Specification

Titer (qPCR analytical results) 3.5E10 TU/mL $\geq 1.0*10^9$

TU: Tranducing Units

Fluorescence detection

(fluorescence expression validation)

Toulouse, the 05 of July, 2019

Anaïs COUFFIN Quality Control Officer

STORAGE: -80°C

The viral vectors must be kept at -80°C immediately upon receipt.

SHELF Life: 1 year from date of receipt under proper storage conditions.

THAWING PROTOCOL

Immediately upon removal from -80°C storage, the vials must be placed directly on ice. Once thawed on ice, place the vials at room temperature just before cell transduction.

VECTOR DILUTION

To dilute the vectors, use a cell culture medium or PBS at room temperature just before cell transduction in order to avoid a thermal shock once the vectors have thawed.



The use of gloves and disposable lab coats when working with viral derived vectors is strongly recommended. This product must be handled only inside a biological safety cabinet with BSL-2 + precautions. The manufacturing of our products is carried out in mycoplasma free cells and in a Biosafety Level 2 facility monitored for microbiological contamination.

The use of lentiviral derived vectors requires laboratory biosafety procedures and practices in accordance with your country regulations:

http://www.cdc.gov/biosafety/publications/bmbl5/index.htm

http://www.hautconseildesbiotechnologies.fr/IMG/pdf/Manuel HCB utilisation confinee OGM.pdf

THIS PRODUCT IS DISTRIBUTED FOR LABORATORY RESEARCH USE ONLY. Your use of this product is also subject to compliance with any applicable licensing requirements described on https://www.takarabio.com/. It is your responsibility to review, understand and adhere to any restrictions imposed by such statements. CAUTION: Not for CLINICAL DIAGNOSTIC OR THERAPEUTIC USE. THE SAFETY AND EFFICACY OF THIS PRODUCT IN DIAGNOSTIC OR OTHER CLINICAL USES HAS NOT BEEN ESTABLISHED.

SYMBOL:



CLASSIFICATION: Biorisk class 2 and BSL 2 for use