

pEF1 α -DsRed-Monomer-C1 Vector

Table of Contents

Product Information	1
Description	2
Location of Features	3
Additional Information	3
Quality Control Data	4

Catalog No.	Amount	Lot Number
631977	10 μ g	Specified on product label.

Product Information

pEF1 α -DsRed-Monomer-C1 is a mammalian expression vector that constitutively expresses a protein of interest fused to the C-terminus of the red fluorescent protein DsRed-Monomer, even after stable integration of the vector into the host cell genome. Stable, constitutive expression of the fusion protein is driven by the human elongation factor 1 alpha (EF1 α) promoter, allowing the monitoring of a variety of cellular processes (such as differentiation in primary or stem cells) without the transgene silencing associated with CMV promoters. The unmodified vector can be used to express modified DsRed-Monomer in mammalian cells.

Package Contents

- 1 tube of pEF1 α -DsRed-Monomer-C1 Vector (20 μ l/tube)

Storage Conditions

- Store plasmid at -20°C .
- Spin briefly to recover contents.
- Avoid repeated freeze/thaw cycles.

Shelf Life

- 1 year from date of receipt under proper storage conditions.

Storage Buffer

- 10 mM Tris-HCl (pH 8.0), 1 mM EDTA (pH 8.0)

Concentration

- 500 ng/ μ l

Shipping Conditions

- Dry ice (-70°C)

Clontech Laboratories, Inc.

A Takara Bio Company
1290 Terra Bella Avenue, Mountain View, CA 94043, USA
U.S. Technical Support: tech@clontech.com

United States/Canada	Asia Pacific	Europe	Japan
800.662.2566	+1.650.919.7300	+33.(0)1.3904.6880	+81.(0)77.543.6116
(PA124335)			




Figure 1. pEF1 α -DsRed-Monomer-C1 vector map. Please note that the vector DNA provided by Clontech is methylated. If you wish to digest the vector with methylation-sensitive enzymes, you will first need to transform the vector into a dam⁻ host strain and purify fresh plasmid DNA.




Figure 2. pEF1 α -DsRed-Monomer-C1 multiple cloning site (MCS).

Description

pEF1 α -DsRed-Monomer-C1 is designed to express a protein of interest fused to the C-terminus of DsRed-Monomer, a monomeric mutant of the *Discosoma sp.* red fluorescent protein DsRed (1). The DsRed-Monomer coding sequence has been human-codon-optimized for high expression in mammalian cells (2). The excitation and emission maxima of native DsRed-Monomer are 557 nm and 585 nm, respectively. Expression of fusion proteins that retain the fluorescence properties of unmodified DsRed-Monomer can be monitored by flow cytometry and localized by fluorescence microscopy.

The multiple cloning site (MCS) in pEF1 α -DsRed-Monomer-C1 is positioned downstream of the DsRed-Monomer coding sequence. Expression of the fusion protein is driven by the EF1 α promoter ($P_{EF1\alpha}$), which remains constitutively active even after stable integration of the vector into the host cell genome (3). A Kozak consensus sequence located immediately upstream of the DsRed-Monomer gene enhances the translational efficiency of the fusion in eukaryotic systems (4), and SV40 polyadenylation signals downstream of the DsRed-Monomer gene direct proper processing of the 3' end of the mRNA.

The vector backbone contains an SV40 origin for replication in mammalian cells expressing the SV40 large T antigen, a pUC origin of replication for propagation in *E. coli*, and an f1 origin for single-stranded DNA production. A neomycin-resistance cassette (Neo^r) allows stably transfected eukaryotic cells to be selected using G418 (5). This cassette consists of the SV40 early promoter (P_{SV40e}), the Tn5 neomycin/kanamycin resistance gene, and polyadenylation signals from the herpes simplex virus thymidine kinase (HSV TK) gene. A bacterial promoter upstream of the cassette drives expression of the kanamycin resistance gene in *E. coli*.

Location of Features

- $P_{EF1\alpha}$ (human elongation factor 1 alpha promoter): 12–1346
- Kozak consensus sequence: 1363–1373
- DsRed-Monomer (human-codon-optimized): 1370–2044
- MCS (multiple cloning site): 2045–2110
- SV40 polyA signal: 2265–2299
- f1 origin of replication: 2362–2817 (complementary)
- P_{SV40e} (SV40 early promoter and enhancer sequences): 2991–3259
- SV40 origin of replication: 3158–3296
- Kan^r/Neo^r (kanamycin/neomycin resistance gene): 3342–4136
- HSV TK polyA signals: 4372–4390
- pUC origin of replication: 4721–5364

Additional Information

Genes cloned into the MCS must be in-frame with the DsRed-Monomer coding sequence, and do not require start or stop codons. The pEF1 α -DsRed-Monomer-C1 vector can be transfected into mammalian cells using any standard transfection method. Cells expressing DsRed-Monomer fusions can be detected by flow cytometry or fluorescence microscopy 12–16 hr after transfection. If required, stable transfectants can be selected using G418 (5). pEF1 α -DsRed-Monomer-C1 can also be used as a cotransfection marker, as the unmodified vector will express DsRed-Monomer in mammalian cells.

Propagation in *E. coli*

- Suitable host strains: DH5 α , HB101 and other general purpose strains. Single-stranded DNA production requires a host containing an F plasmid, such as the JM109 or XL1-Blue strains.
- Selectable marker: plasmid confers resistance to kanamycin (50 μ g/ml) in *E. coli* hosts.
- *E. coli* replication origin: pUC
- Copy number: high

Excitation and Emission Maxima of DsRed-Monomer

- Excitation: 557 nm
- Emission: 585 nm

References

1. Matz, M. V., *et al.* (1999) *Nat. Biotechnol.* **17**(10):969–973.
2. Haas, J., *et al.* (1996) *Curr. Biol.* **6**(3):315–324.
3. Wang, R. *et al.* (2008) *Stem Cells Dev.* **17**(2):279–289.
4. Kozak, M. (1987) *Nucleic Acids Res.* **15**(20):8125–8148.
5. Gorman, C. (1985) In *DNA Cloning: A Practical Approach, Vol. II*. Ed. D. M. Glover (IRL Press, Oxford, U.K.) pp. 143–190.

Quality Control Data

Plasmid Identity & Purity

- Digestion with the indicated restriction enzymes produced fragments of the indicated sizes on a 0.8% agarose/EtBr gel:

Enzyme(s)	Fragment(s)
BamHI	5.4 kb
AgeI	1.1 & 4.3 kb

- Vector identity was confirmed by sequencing.
- A_{260}/A_{280} : 1.8–2.0

pEF1alpha-DsRed-Monomer-C1 Vector

CATALOG NO.

631977

NOTICE TO PURCHASER:

Our products are to be used for **Research Use Only**. They may not be used for any other purpose, including, but not limited to, use in humans, therapeutic or diagnostic use, or commercial use of any kind. Our products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products or to provide a service to third parties without our prior written approval.

Your use of this product is also subject to compliance with the licensing requirements, listed below if applicable, and described on the product's web page at <http://www.takarabio.com>. It is your responsibility to review, understand and adhere to any restrictions imposed by these statements.

TRADEMARKS:

©2026 Takara Bio Inc. All Rights Reserved.

All trademarks are the property of Takara Bio Inc. or its affiliate(s) in the U.S. and/or other countries or their respective owners. Certain trademarks may not be registered in all jurisdictions.

Takara Bio USA, Inc.

2560 Orchard Parkway, San Jose, CA 95131, USA
U.S. Technical Support: technical_support@takarabio.com

United States/Canada

800.662.2566

Asia Pacific

+1.650.919.7300

Europe

+33.(0)1.3904.6880

Japan

+81.(0)77.565.6999

3/13/2026