# Certificate of Analysis



## Advantage® GC Genomic LA Polymerase Mix

Catalog No. Amount **Lot Number** 

639153 200 reactions Specified on product label.

## **Description**

The Advantage GC Genomic LA Polymerase Mix is composed of a full-length Taq DNA polymerase, a small amount of proofreading enzyme, and a hot-start antibody that inhibits polymerase activity at ambient temperature. The 2X Advantage GC-Melt Buffer is specifically designed to amplify DNA templates with high GC content or a significant amount of secondary structure. This combination allows synthesis of products up to 20 kb for complex templates, such as genomic DNA containing GC-rich regions, and up to 48 kb for less complex templates, such as λ DNA. The polymerase mix has 6.5X higher fidelity than wild-type Taq DNA polymerase, due to the presence of the proofreading 3'-to-5' exonuclease. The robust enzyme/buffer system requires less optimization and produces higher yields than other "long and accurate polymerases." Sufficient polymerase mix and buffer are provided for 200 reactions (25 µl each).

### **Package Contents**

- 50 μl Advantage GC Genomic LA Polymerase (5 units/μl)
- 2 x 1.25 ml 2X Advantage GC-Melt Buffer (containing 5 mM MgCl<sub>2</sub>)

## **Storage Conditions**

Store at -20°C

#### Shelf Life

1 year from date of receipt under proper storage conditions.

### **Shipping Conditions**

Dry ice  $(-70^{\circ}C)$ 

#### **Product Documents**

Documents for our products are available for download at takarabio.com/manuals The following documents apply to this product:

Advantage GC Genomic LA Polymerase Mix Protocol-at-a-Glance (PT3938-2)

## **Quality Control Data**

### **Raw Material Quality Control**

#### **Purity**

No nicking activity, endonuclease activity, or exonuclease activity was detected after incubation of 0.6 mg of supercoiled pBR322 DNA, 0.65 mg of  $\lambda$  DNA, or 0.6 mg of  $\lambda$ -Hind III digest with 10 units of this enzyme for 1 hour at 74°C.

Advantage GC Genomic LA Polymerase Mix

### **Unit definition**

One unit is the amount of enzyme that will incorporate 10 nmol of dNTP into acid-insoluble products in 30 minutes at 74°C using activated salmon sperm DNA as template.

#### Reaction mixture for unit definition

25 mM TAPS (pH 9.3 at 25°C)
50 mM KCl
2 mM MgCl<sub>2</sub>
1 mM 2-mercaptoethanol
200 mM Each dATP, dGTP, dTTP
100 mM [α-<sup>32</sup>P]-dCTP
0.25 mg/ml activated salmon sperm DNA

## **Functional Quality Control**

## **PCR** performance

The Advantage GC Genomic LA Polymerase Mix was tested for PCR performance using a  $\lambda$  DNA control template and primers. Reactions (50  $\mu$ l) were assembled and performed using a standard PCR protocol. When the PCR products were examined by electrophoresis on a 1.0% agarose/EtBr gel, a major band of 35 kb was observed.

To ensure optimum PCR performance, the Advantage GC Genomic LA Polymerase Mix was also tested using human genomic DNA as template. Reactions were assembled and performed using a standard PCR protocol. When the PCR products were examined by electrophoresis on a 1.0% agarose/EtBr gel, a major band of 17.5 kb was observed.

To ensure optimal performance of the GC-Melt Buffer, a GC-rich fragment (65% GC content) from the *c-jun* gene was amplified using human genomic DNA as template. Reactions were assembled using either the 2X Advantage GC-Melt Buffer or a standard buffer, and performed using a standard PCR protocol. When the PCR products were examined by electrophoresis on a 1.0% agarose/EtBr gel, a major 1.2-kb band was observed in reactions containing the GC-Melt Buffer. The improvement (no non-specific band) was observed in GC-Melt Buffer by comparison with LA Buffer.

## **Hot-start** antibody

Hot-start antibody inhibition of the Advantage GC Genomic LA enzyme was confirmed to be greater than 80% following a 10 min incubation of the reaction at 55°C.

It is certified that this product meets the above specifications, as reviewed and approved by the Quality Department.

(101717) Page 2 of 2



## Advantage® GC Genomic LA Polymerase Mix

## CATALOG NO.

639153

#### NOTICE TO PURCHASER:

Our products are to be used for research purposes only. They may not be used for any other purpose, including, but not limited to, use in drugs, in vitro diagnostic purposes, therapeutics, or in humans. Our products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products or to provide a service to third parties without prior written approval of Takara Bio USA, Inc.

Your use of this product is also subject to compliance with the licensing requirements listed below and described on the product's web page at <a href="http://www.takarabio.com">http://www.takarabio.com</a>. It is your responsibility to review, understand and adhere to any restrictions imposed by these statements.

#### TRADEMARKS:

#### ©2017 Takara Bio Inc. All Rights Reserved.

All trademarks are the property of Takara Bio Inc. or its affiliate(s) in the U.S. and/or other countries or their respective owners. Certain trademarks may not be registered in all jurisdictions.

Takara Bio USA, Inc.

1290 Terra Bella Avenue, Mountain View, CA 94043, USA

U.S. Technical Support: techUS@takarabio.com
United States/Canada Asia Pacific

800.662.2566 +1.650.919.7300 +33.(0)1.3904.6880 +81.(0)77.565.6999

**Europe** 

Japan