


Keeping up with the microbes: Methods for effective AMR detection



Why is AMR detection important?


Antimicrobial resistance (AMR) is an emerging public health crisis in which microorganisms—such as bacteria, fungi, viruses, and parasites—acquire antibiotic resistance genes (ARGs) that enable them to withstand antimicrobial drugs. As a result, combatting these microorganisms requires constant AMR surveillance and new antibiotic development.

With AMR on the rise, we need effective methods to identify and limit resistant microbes.



How do scientists detect AMR?

Microorganisms with AMR can be distinguished either by phenotype, such as survival of antibiotic exposure, or molecular indicators, such as expression of ARGs. While AMR detection strategies constantly evolve, there are three primary methods used worldwide:



Bacterial culture plates

Microorganism samples are grown on bacterial culture plates under antibiotic exposure, with surviving microorganisms considered positive for AMR.

- ✓ Simple and straightforward protocol
- ✓ Inexpensive, requiring few materials
- ✗ Long turnaround time (>1 day)
- ✗ Unsuitable for nonculturable microorganisms
- ✗ No quantitative results
- ✗ Labor-intensive



Sequencing

Sequencing identifies ARGs or ARG expression in microorganisms as an indicator of AMR.

- ✓ Covers whole genome or transcriptome
- ✓ Processes multiple samples at once
- ✗ Requires lengthy turnaround time (2–3 days)
- ✗ Involves technical expertise for complex data analysis
- ✗ No quantitative results




Plate-based qPCR

Plate-based qPCR (with 96 or 384 wells) detects the presence of specific ARG targets in sample microorganisms.


- ✓ Fast turnaround time (~3 hr)
- ✓ Minimal expertise required
- ✓ Quantitative, reliable results
- ✓ Inexpensive at small scale
- ✗ Limited throughput (max 4 samples per plate for traditional 96-target panels)
- ✗ Difficult to scale up, due to high reagent costs and labor needs

As demand grows for routine, large-scale AMR monitoring of hundreds of ARG targets, it is evident that these traditional methods fall short for global AMR surveillance efforts.

What does effective large-scale AMR look like?



SmartChip ND Real-Time PCR System



References

- Antimicrobial Resistance Facts and Stats | Antimicrobial Resistance | CDC. at <<https://www.cdc.gov/antimicrobial-resistance/data-research/facts-stats/index.html>>
- Economou, V. & Gousia, P. Agriculture and food animals as a source of antimicrobial-resistant bacteria. *Infect. Drug Resist.* 49–61 (2015).
- Multi-drug resistant gonorrhoea. at <<https://www.who.int/news-room/fact-sheets/detail/multi-drug-resistant-gonorrhoea>>
- Naghavi, M. et al. Global burden of bacterial antimicrobial resistance 1990–2021: a systematic analysis with forecasts to 2050. *Lancet* **404**, 1199–1226 (2024).
- Raccagni, A. R. et al. Neisseria gonorrhoeae antimicrobial resistance: the future of antibiotic therapy. *J. Clin. Med.* **12**, 7767 (2023).
- Uddin, T. M. et al. Antibiotic resistance in microbes: History, mechanisms, therapeutic strategies and future prospects. *J. Infect. Public Health* **14**, 1750–1766 (2021).



Learn more about high-throughput AMR detection:
takarabio.com/amr-detection

that's
GOOD
 science!®

Takara Bio USA, Inc.

United States/Canada: +1.800.662.2566 • Asia Pacific: +1.650.919.7300 • Europe: +33.(0)1.3904.6880 • Japan: +81.(0)77.565.6999

For Research Use Only. Not for use in diagnostic procedures.

© 2025 Takara Bio Inc. All Rights Reserved. All trademarks are the property of Takara Bio Inc. or its affiliate(s) in the U.S. and/or other countries or their respective owners. Certain trademarks may not be registered in all jurisdictions. Additional product, intellectual property, and restricted use information is available at takarabio.com.

6.25 US (634190)